

Moment varieties for mixtures of products

Yulia Alexandr (UC Berkeley)
joint work with Joe Kileel and Bernd Sturmfels

CSU Fresno
May 6, 2023

The (nonparametric) set-up

Consider n independent random variables X_1, X_2, \dots, X_n on the line \mathbb{R} .

Assumptions:

- ★ **No** assumptions about X_k , only that moments $\mu_{ki} = \mathbb{E}(X_k^i)$ exist.
- ★ The moments μ_{ki} are unknowns.
- ★ The only equations we require are $\mu_{k0} = 1$ for $k = 1, 2, \dots, n$.

We consider a random variable X on \mathbb{R}^n that is the product of these n arbitrary independent random variables on \mathbb{R} . By independence, we have

$$\mathbb{E}(X_1^{i_1} X_2^{i_2} \cdots X_n^{i_n}) = \mathbb{E}(X_1^{i_1}) \cdot \mathbb{E}(X_2^{i_2}) \cdots \mathbb{E}(X_n^{i_n}).$$

This leads us to the *moment variety* $\mathcal{M}_{n,d}$, which has parametrization

$$m_{i_1 i_2 \dots i_n} = \mu_{1 i_1} \mu_{2 i_2} \cdots \mu_{n i_n} \text{ where } i_1, i_2, \dots, i_n \geq 0 \text{ and } i_1 + i_2 + \cdots + i_n = d.$$

The image is a toric variety of dimension at most $nd - 1$ in $\mathbb{P}^{\binom{n+d-1}{d}-1}$.

Example

Consider $\mathcal{M}_{5,3}$ in \mathbb{P}^{34} . The solutions to $i_1 + i_2 + i_3 + i_4 + i_5 = 3$ can be grouped into three partitions: $\lambda = (1\ 1\ 1)$, $\lambda = (2\ 1)$, $\lambda = (3)$. Consider the following three toric varieties of dimensions 4, 8, 4 respectively:

$$\mathcal{M}_{5,(111)} \subset \mathbb{P}^9 : m_{11100} = \mu_{11}\mu_{21}\mu_{31}, \dots, m_{00111} = \mu_{31}\mu_{41}\mu_{51}.$$

$$\mathcal{M}_{5,(21)} \subset \mathbb{P}^{19} : m_{21000} = \mu_{12}\mu_{21}, m_{12000} = \mu_{11}\mu_{22}, \dots, m_{00012} = \mu_{41}\mu_{52}.$$

$$\mathcal{M}_{5,(3)} = \mathbb{P}^4 : m_{30000} = \mu_{13}, m_{03000} = \mu_{23}, \dots, m_{00003} = \mu_{53}.$$

Combining these parametrizations yields the original variety.

We will also study $\mathcal{M}_{n,d}$ under projections $\mathbb{P}^{\binom{n+d-1}{d}-1} \dashrightarrow \mathbb{P}^{|\mathcal{N}_\lambda|-1}$ for any partition λ of d with $\leq n$ parts. We denote these toric varieties by $\mathcal{M}_{n,\lambda}$.

Toric combinatorics

First, we are interested in studying the toric varieties $\mathcal{M}_{n,d}$ and $\mathcal{M}_{n,\lambda}$.

Familiar examples:

- ★ For any n , consider the partition $\lambda = (1^d) = (1 \ 1 \dots 1)$ of $d < n$. Then $\mathcal{M}_{n,(1^d)}$ is the associated toric variety to the **hypersimplex**

$$\Delta(n, d) = \text{conv}\{e_{\ell_1} + e_{\ell_2} + \dots + e_{\ell_d} : 1 \leq \ell_1 < \ell_2 < \dots < \ell_d \leq n\}$$

It has dimension $n - 1$ in $\mathbb{P}^{\binom{n}{d}-1}$.

- ★ Consider the partition $\lambda = (n - 1, n - 2, \dots, 2, 1)$. Then the moment variety $\mathcal{M}_{n,\lambda}$ is the toric variety of the **Birkhoff polytope**, which lives in $\mathbb{P}^{n!-1}$ and has dimension $(n - 1)^2$.

Toric results

Theorem (A., Kileel, Sturmfels)

The dimension of the moment variety $\mathcal{M}_{n,d}$ is $\min \left\{ nd - 1, \binom{n+d-1}{d} - 1 \right\}$.

Given a partition λ of length n , let $k_0 \geq k_1 \geq \dots \geq k_s$ be multiplicities of the distinct parts in λ . We define

$$\nu = (\underbrace{s, \dots, s}_{k_s}, \underbrace{s-1, \dots, s-1}_{k_{s-1}}, \dots, \underbrace{1, \dots, 1}_{k_1}, \underbrace{0, \dots, 0}_{k_0}).$$

to be the *reduction* of λ . Here $s+1$ is the number of distinct parts of λ .

Ex: both $(8, 5, 5, 4)$ and $(7, 7, 3, 0)$ reduce to $\nu = (2, 1, 0, 0)$, with $s = 2$.

Ex: if $\lambda = (1^d)$, we recover the identification $\Delta(n, d)$ with $\Delta(n, n-d)$.

Theorem (A., Kileel, Sturmfels)

The moment variety $\mathcal{M}_{n,\lambda} = \mathcal{M}_{n,\nu}$ has dimension $(n-1)s$.

What about generators?

Example

Consider the variety $\mathcal{M}_{4,4}$ in \mathbb{P}^{34} . Its ideal is generated by 52 quadrics and 28 cubics. The subset of the generators which involves the twelve unknowns $m_{2110}, \dots, m_{0112}$ does not suffice to cut out $\mathcal{M}_{4,(211)}$ in \mathbb{P}^{11} .

Theorem (A., Kileel, Sturmfels)

For any partition λ , the ideal of $\mathcal{M}_{n,\lambda}$ is generated by quadrics and cubics.

The ideals for $\mathcal{M}_{n,d}$ are more complicated. We conjecture that there does not exist a uniform degree bound for their generators.

Mixtures

Now we consider the mixtures of r copies of our toric models.

Algebraically, these are the *secant varieties* $\sigma_r(\mathcal{M}_{n,d})$ and $\sigma_r(\mathcal{M}_{n,\lambda})$. The first is parametrized by

$$m_{i_1 i_2 \dots i_n} = \sum_{j=1}^r \mu_{1i_1}^{(j)} \mu_{2i_2}^{(j)} \cdots \mu_{ni_n}^{(j)} \text{ with } i_1, i_2, \dots, i_n \geq 0 \text{ and } i_1 + i_2 + \dots + i_n = d.$$

These varieties are no longer toric! What can we say about their dimensions, degrees, generators?

◇ Consider the secant variety $\sigma_2(\mathcal{M}_{5,2})$. The parametrization is given as

$$m_{20000} = \mu_{12}^{(1)} + \mu_{12}^{(2)} \quad , \quad \dots \quad , \quad m_{11000} = \mu_{11}^{(1)} \mu_{21}^{(1)} + \mu_{11}^{(2)} \mu_{21}^{(2)} \quad , \quad \dots$$

Example (continued)

Note $\mathcal{M}_{5,2} = \mathcal{M}_{5,(2)} \star \mathcal{M}_{5,(11)} = \mathbb{P}^4 \star \mathcal{M}_{5,(11)}$, since


$$\begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} & m_{15} \\ m_{12} & m_{22} & m_{23} & m_{24} & m_{25} \\ m_{13} & m_{23} & m_{33} & m_{34} & m_{35} \\ m_{14} & m_{24} & m_{34} & m_{44} & m_{45} \\ m_{15} & m_{25} & m_{35} & m_{45} & m_{55} \end{bmatrix} = \begin{bmatrix} \mu_{12} & \mu_{11}\mu_{21} & \mu_{11}\mu_{31} & \mu_{11}\mu_{41} & \mu_{11}\mu_{51} \\ \mu_{11}\mu_{21} & \mu_{22} & \mu_{21}\mu_{31} & \mu_{21}\mu_{41} & \mu_{21}\mu_{51} \\ \mu_{11}\mu_{31} & \mu_{21}\mu_{31} & \mu_{32} & \mu_{31}\mu_{41} & \mu_{31}\mu_{51} \\ \mu_{11}\mu_{41} & \mu_{21}\mu_{41} & \mu_{31}\mu_{41} & \mu_{42} & \mu_{41}\mu_{51} \\ \mu_{11}\mu_{51} & \mu_{21}\mu_{51} & \mu_{31}\mu_{51} & \mu_{41}\mu_{51} & \mu_{52} \end{bmatrix}.$$

$$\sigma_2(\mathcal{M}_{5,2}) = \sigma_2(\mathbb{P}^4 \star \mathcal{M}_{5,(11)}) = \mathbb{P}^4 \star \boxed{\sigma_2(\mathcal{M}_{5,(11)})} \subset \mathbb{P}^4 \star \mathbb{P}^9 = \mathbb{P}^{14}.$$

The ideal of $\sigma_2(\mathcal{M}_{5,(11)})$ is principal, generated by the pentad

$$\begin{aligned} & m_{12}m_{13}m_{24}m_{35}m_{45} - m_{12}m_{13}m_{25}m_{34}m_{45} - m_{12}m_{14}m_{23}m_{35}m_{45} + m_{12}m_{14}m_{25}m_{34}m_{35} \\ & + m_{12}m_{15}m_{23}m_{34}m_{45} - m_{12}m_{15}m_{24}m_{34}m_{35} + m_{13}m_{14}m_{23}m_{25}m_{45} - m_{13}m_{14}m_{24}m_{25}m_{35} \\ & - m_{13}m_{15}m_{23}m_{24}m_{45} + m_{13}m_{15}m_{24}m_{25}m_{34} + m_{14}m_{15}m_{23}m_{24}m_{35} - m_{14}m_{15}m_{23}m_{25}m_{34}. \end{aligned}$$

This is the factor analysis
model $\mathbf{F}_{5,2}$.



Dimensions of mixtures

Proposition (A., Kileel, Sturmfels)

The dimension of the moment variety satisfies the upper bound

$$\dim(\sigma_r(\mathcal{M}_{n,d})) \leq \min\{rnd - rn + n - 1, \binom{n+d-1}{d} - 1\}. \quad (1)$$

Consider $\sigma_2(\mathcal{M}_{5,3}) = \sigma_2(\mathcal{M}_{5,(3)} \star \widetilde{\mathcal{M}}_{5,3}) = \mathbb{P}^4 \star \sigma_2(\widetilde{\mathcal{M}}_{5,3})$ in \mathbb{P}^{34} . Note:

- ★ we know $\dim(\widetilde{\mathcal{M}}_{5,3}) \leq (5 \cdot 3 - 5) - 1 = 9$,
- ★ so $\dim(\sigma_2(\mathcal{M}_{5,3})) \leq 4 + 1 + (2 \cdot 9 + 1) = 24$.

Theorem (A., Kileel, Sturmfels)

The dimension $\sigma_r(\mathcal{M}_{n,d})$ is bounded above by the optimal value of

maximize $c_1 + c_2 + \dots + c_d - 1$	subject to	$0 \leq c_i \leq nr$ for $i \in [d]$
	and	$\sum_{i \in S} c_i \leq \sum_{\lambda \cap S \neq \emptyset} N_\lambda $ for $S \subseteq [d]$.

The last sum ranges over partitions $\lambda \vdash d$ of length $\leq n$ having nonempty intersection with S .

Conjecture: this bound is tight for $d \geq 3$!

Implicitization

Solving the implicitization problem is difficult!

Remark: Our initial parametrization is not one-to-one. If ω is a primitive d th root of unity then we can replace μ_{ki} by $\mu_{ki} \omega^i$ without changing $m_{i_1 i_2 \dots i_n}$. We parameterize to make degree computations in **Julia** faster.

Consider the variety $\mathcal{M}_{6,(111)}$. Its ideal is given by the 2×2 minors of

$$\begin{bmatrix} \star & \star & \star & \star & \star & m_{123} & m_{124} & m_{125} & m_{126} & m_{134} & m_{135} & m_{136} & m_{145} & m_{146} & m_{156} \\ \star & m_{123} & m_{124} & m_{125} & m_{126} & \star & \star & \star & \star & m_{234} & m_{235} & m_{236} & m_{245} & m_{246} & m_{256} \\ m_{123} & \star & m_{134} & m_{135} & m_{136} & \star & m_{234} & m_{235} & m_{236} & \star & \star & \star & m_{345} & m_{346} & m_{356} \\ m_{124} & m_{134} & \star & m_{145} & m_{146} & m_{234} & \star & m_{245} & m_{246} & \star & m_{345} & m_{346} & \star & \star & m_{456} \\ m_{125} & m_{135} & m_{145} & \star & m_{156} & m_{235} & m_{245} & \star & m_{256} & m_{345} & \star & m_{356} & \star & m_{456} & \star \\ m_{126} & m_{136} & m_{146} & m_{156} & \star & m_{236} & m_{246} & m_{256} & \star & m_{346} & m_{356} & \star & m_{456} & \star & \star \end{bmatrix}$$

The ideal of $\sigma_2(\mathcal{M}_{6,(111)})$ is generated by 20 cubics and 12 quintics. The ideal of $\sigma_3(\mathcal{M}_{6,(111)})$ has no quadrics or cubics, but contains a unique quartic. Computations in **Julia** reveal:

$$\deg(\sigma_2(\mathcal{M}_{6,(111)})) = 465 \quad \text{and} \quad \deg(\sigma_3(\mathcal{M}_{6,(111)})) = 80.$$

More implicitization

Proposition (A., Kileel, Sturmfels)

The variety $\sigma_2(\mathcal{M}_{5,3})$ has dimension 24 and degree 3225 in \mathbb{P}^{34} . Its prime ideal is generated by 313 polynomials, namely 10 cubics, 283 quintics, 10 sextics and 10 septics. These are obtained by elimination from the ideal of 3×3 minors of the 5×15 matrix

$$\begin{bmatrix} a_{23} & a_{24} & a_{25} & a_{34} & a_{35} & a_{45} & \star & \star & \star & \star & \star & b_{21} & b_{31} & b_{41} & b_{51} \\ a_{13} & a_{14} & a_{15} & \star & \star & \star & a_{34} & a_{35} & a_{45} & \star & b_{12} & \star & b_{32} & b_{42} & b_{52} \\ a_{12} & \star & \star & a_{14} & a_{15} & \star & a_{24} & a_{25} & \star & a_{45} & b_{13} & b_{23} & \star & b_{43} & b_{53} \\ \star & a_{12} & \star & a_{13} & \star & a_{15} & a_{23} & \star & a_{25} & a_{35} & b_{14} & b_{24} & b_{34} & \star & b_{54} \\ \star & \star & a_{12} & \star & a_{13} & a_{14} & \star & a_{23} & a_{24} & a_{34} & b_{15} & b_{25} & b_{35} & b_{45} & \star \end{bmatrix}.$$

Proposition (A., Kileel, Sturmfels)

The variety $\sigma_2(\mathcal{M}_{4,4})$ has dimension 27 and degree 8650 in \mathbb{P}^{34} . Its prime ideal has only three minimal generators in degrees at most six.

Finiteness up to symmetry

Our ideals satisfy natural inclusions

$$I(\sigma_r(\mathcal{M}_{n,\bullet})) \subset I(\sigma_r(\mathcal{M}_{n+1,\bullet})), \quad \text{where } \bullet \in \{d, \lambda\},$$

by appending a zero to the indices of every coordinate: $m_{i_1 i_2 \dots i_n} \mapsto m_{i_1 i_2 \dots i_n 0}$. Iterate these inclusions and let the big symmetric group act:

$$\langle S_n I(\sigma_r(\mathcal{M}_{n_0,\bullet})) \rangle \subseteq I(\sigma_r(\mathcal{M}_{n,\bullet})) \quad \text{for } n > n_0.$$

Ideal-theoretic finiteness means $\exists n_0$ such that equality holds for $n > n_0$.

Theorem (A., Kileel, Sturmfels)

Given any partition $\lambda \vdash d$ and integer $r \geq 1$, set-theoretic finiteness holds for the varieties $\sigma_r(\mathcal{M}_{n,d})$ and $\sigma_r(\mathcal{M}_{n,\lambda})$. Ideal-theoretic finiteness holds in the toric case $r = 1$.

Example

The ideal of the variety $\mathcal{M}_{n,(1^d)}$ is generated by quadrics. The indices occurring in each quadratic binomial are 1 in at most $2d$ of the n coordinates. Therefore, ideal-theoretic finiteness holds with $n_0 = 2d$.

If $\lambda = (1\ 1)$, then $n_0 = 4$. Indeed:

$$I(\mathcal{M}_{4,\lambda}) = \langle m_{0101}m_{1010} - m_{0110}m_{1001}, m_{0011}m_{1100} - m_{0110}m_{1001} \rangle$$

$$\begin{aligned} I(\mathcal{M}_{5,\lambda}) = \langle &m_{01001}m_{10100} - m_{01100}m_{10001}, m_{00011}m_{10100} - m_{00110}m_{10001}, \\ &m_{11000}m_{00101} - m_{01100}m_{10001}, m_{10010}m_{00101} - m_{00110}m_{10001}, \\ &m_{10010}m_{01100} - m_{10100}m_{01010}, m_{00011}m_{01100} - m_{00101}m_{01010}, \\ &m_{00110}m_{11000} - m_{10100}m_{01010}, m_{00011}m_{11000} - m_{10001}m_{01010}, \\ &m_{01001}m_{10010} - m_{10001}m_{01010}, m_{00110}m_{01001} - m_{00101}m_{01010} \rangle \end{aligned}$$

Corollary

Fix a partition λ with e nonzero parts, and suppose that n increases. The toric varieties $\mathcal{M}_{n,\lambda}$ satisfy ideal-theoretic finiteness for some $n_0 \leq 3e$.

Thanks!